

VLT AND HST OBSERVATIONS OF THE HOST GALAXY OF GRB 990705

E. LE FLOC'H³, P.-A. DUC³, I.F. MIRABEL^{3,4}, D.B. SANDERS^{5,6}, G. BOSCH⁷, I. RODRIGUES³,
 T.J.-L. COURVOISIER^{8,9}, S. MEREGHETTI¹⁰, & J. MELNICK¹¹

Received 2002 September 6; Accepted 2002 November 12

ABSTRACT

We present VLT spectroscopic observations of Gamma-Ray Burst (GRB) 990705 host galaxy and highlight the benefits provided by the prompt-phase features of GRBs to derive the redshifts of the latter. In the host spectrum, we indeed detect an emission feature which we attribute to the [OII] $\lambda\lambda$ 3726/3729 Å doublet and derive an unambiguous redshift $z=0.8424 \pm 0.0002$ for this galaxy. This is in full agreement with the value $z \sim 0.86 \pm 0.17$ previously derived using a transient absorption edge discovered in the X-ray spectrum of GRB 990705. This burst is therefore the first GRB for which a reliable redshift was derived *from the prompt phase emission itself*, as opposed to redshift determinations performed using putative host galaxy emission lines or interstellar absorption lines in the GRB afterglows. Deep and high resolution images of the host of GRB 990705 with the HST/STIS camera reveal that the burst occurred in a nearly face-on Sc spiral galaxy typical of disk-dominated systems at $0.75 \leq z \leq 1$. Assuming a cosmology with $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$, we derive an absolute B magnitude $M_B = -21.75$ for this galaxy and a star formation rate $\text{SFR} \approx 5\text{--}8 \text{ M}_\odot \text{ yr}^{-1}$. Finally, we discuss the implications of using X-ray transient features to derive GRB redshifts with larger burst samples, and especially examine the case of short and dark-long GRBs.

Subject headings: galaxies: spiral — galaxies: starburst — galaxies: individual (GRB 990705 host) — gamma rays: bursts

1. INTRODUCTION

Since the discovery of their X-ray, optical and radio transient counterparts, the cosmic Gamma-ray bursts (GRBs) have been regarded as one of the most promising tools to probe the star formation in the early Universe (Totani 1997; Wijers et al. 1998; Mirabel et al. 2000; Blain & Natarajan 2000). There is indeed increasing evidence that the long and soft GRBs originate from the core collapse of massive stars within starburst regions of distant galaxies (e.g., Bloom02a). Since they are likely detectable up to very high redshifts (Lamb & Reichart 2000), GRBs could soon open a new window to sample the star-forming activity at cosmological lookback times, and ultimately provide a new glimpse on galaxy evolution.

The possibility to detect emission and/or absorption features in the spectra of GRBs and their afterglows is among the most outstanding benefits of the high- z galaxy selection by these events. Such detections can indeed be done independently of the GRB host luminosities and have already enabled spectroscopic redshifts of very faint galaxies to be derived (e.g., Vreeswijk01). This perspective strongly contrasts with the deep survey observations which can only provide photometric redshifts for

the faintest sources. Nonetheless, the correct GRB redshift identifications from the lines detected in *afterglow* spectra are not always straightforward. Absorption features observed in the optical continuum of GRB counterparts may indeed originate from foreground absorbers (e.g., Metzger97), while the interpretation of the emission lines detected in the X-ray afterglows has already led to some mis-identifications of host redshifts. (e.g., GRB 970828; Yoshida99, Djorgovski01a).

In this context, the GRB 990705 event is of remarkable interest. Using data from the BeppoSAX satellite, Amati et al. (2000) reported the discovery of a transient absorption edge at $\sim 3.8 \text{ keV}$ in the prompt X-ray emission of this burst, which they interpreted as the GRB-intrinsic signature of an iron-enriched absorbing medium at $z \sim 0.86$ (see also Lazzati01). This has been so far the only GRB for which a feature allowing a possible redshift determination was observed during the *prompt emission* of the burst itself. Following this event, an optical and near-infrared follow-up by Masetti et al. (2000) led to the discovery of a red and rapidly-decaying afterglow localized behind the Large Magellanic Cloud (LMC), while deep optical images of the burst location performed with the HST (Holland

¹ Based on observations with the Very Large Telescope, obtained at the European Southern Observatory in Chile under Prop. 68.B-0250(B)
² Based on observations with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc. under NASA contract No. NAS5-26555.
³ CEA/DSM/DAPNIA Service d'Astrophysique, F-91191 Gif-sur-Yvette, France (elefloch, paduc, fmirabel, irapuan@cea.fr)
⁴ Instituto de Astronomía y Física del Espacio, cc 67, suc 28. 1428 Buenos Aires, Argentina
⁵ Institute for Astronomy, University of Hawai'i, 2680 Woodlawn Drive, Honolulu, HI 96822, United States (sanders@ifa.hawaii.edu)
⁶ Max-Planck Institute for Extraterrestrial Physik, D-85740, Garching, Germany
⁷ Facultad de Cs. Astronomicas y Geofisica, Paseo del Bosque s/n, La Plata, Argentina (guille@lahuan.fcaglp.unlp.edu.ar)
⁸ INTEGRAL Science Data center, Ch. d'Ecogia 16, CH-1290 Versoix, Switzerland (Thierry.Courvoisier@obs.unige.ch)
⁹ Geneva Observatory, Ch. des Maillettes 11, 1290 Sauverny, Switzerland
¹⁰ Istituto di Astrofisica Spaziale e Fisica Cosmica, Sezione di Milano "G. Occhialini," via Bassini 15, I-20133 Milan, Italy (sandro@mi.iasf.cnr.it)
¹¹ European Southern Observatory, Alonso de Cordova 3107, Santiago, Chile (jmelnick@eso.org)

et al. 2000a) and the VLT (Saracco et al. 2001) revealed an $R \sim 22.2$ mag underlying host galaxy.

In this letter we report on VLT observations carried out to derive the spectroscopic redshift of this host galaxy, and which allowed us to confirm the redshift of GRB 990705 derived by Amati et al. (2000). We also analyse public HST data of the host.

2. OBSERVATIONS AND DATA REDUCTION

The spectroscopic observations of the GRB 990705 host galaxy were performed on 2001 December 21 and 22 (burst trigger + ~ 900 days) with the FORS2 instrument installed on the VLT UT4/Yepun at ESO. Spectra were obtained under moderate seeing conditions ($\sim 1''$) using a medium resolution grism (600RI) in combination with a $1''$ -width slit, and totalizing an integration time of 1.5 hours. We thus covered an effective wavelength range $\sim 5600 - 8000 \text{ \AA}$ with an instrumental resolution $\sim 4.5 \text{ \AA}$. The slit was positioned on the sky so as to cover the outer region of the galaxy where the burst occurred. The galaxy spectra were flux-calibrated using spectroscopic standard stars.

The HST observations of the GRB 990705 host galaxy¹² were taken and reduced by Holland et al. (2000a,b) as part of the ‘‘Survey of the Host Galaxies of Gamma-Ray Bursts’’ using the STIS camera. Images were obtained with the 50CCD (clear, pivot $\lambda_o = 5835 \text{ \AA}$, hereafter CL) and F28X50LP (long pass, pivot $\lambda_o = 7208 \text{ \AA}$, hereafter LP) apertures, respectively on 2000 July 25 and 2000 August 25 (i.e., ~ 400 days after the burst). The respective total exposure times were 8851s and 8202s in the CL and LP apertures. We deconvolved the images following a multi-resolution wavelet decomposition (Starck et al. 1998) and the use of PSFs obtained from the combination of foreground stars in the images.

The photometry measurements were performed on the data before deconvolution to preserve reliable flux and noise estimates. We corrected the CL and LP aperture data from absorptions $A_{CL} = 0.36$ and $A_{LP} = 0.28$ mag assuming the extinction curve of Cardelli et al. (1989) and the Galactic + LMC extinction $E(B - V) = 0.12$ obtained by Dutra et al. (2001). Moreover, we carried out a careful analysis using a multi-resolution transform method to subtract from the images the multiple LMC foreground stars superimposed on the plane of the galaxy.

3. RESULTS

3.1. The redshift of the GRB 990705 host galaxy

The final VLT spectrum is shown in Figure 1. An emission feature is clearly detected at $\sim 6868 \text{ \AA}$ in a region of the spectrum where the residuals from the sky line subtraction are negligible. Attributing this feature respectively to $H\alpha$ or $Ly\alpha$ would imply redshifts $z = 0.05$ and $z = 4.66$, which is inconsistent with the spiral morphology and the angular size of the galaxy (see section 3.2). The line can thus only be due to $[OII] \lambda\lambda 3726/3729 \text{ \AA}$. It is actually not resolved in our spectrum, but its width is in fact consistent with that of the $[OII]$ doublet. Note that the low signal to noise ratio longward of 7300 \AA does not allow us to detect $H\delta$ and $H\gamma$. From the $[OII]$ line, we derive a secure heliocentric redshift $z = 0.8424 \pm 0.0002$

for the host galaxy and GRB 990705. This is in full agreement with the value $z \sim 0.86 \pm 0.17$ obtained by Amati et al. (2000) from the transient feature observed in the GRB prompt emission, and also appears consistent with the redshift $z = 0.843$ already mentioned by Lazati et al. (2001). Assuming a standard cosmology with $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$, we thus measure for the host of GRB 990705 a luminosity distance $d_l = 5.8 \text{ Gpc}$, and a projected scale of 8.2 proper kpc (or 15.2 comoving kpc) per arcsecond on the sky. Because of the extended and rather diffuse emission of the galaxy (see section 3.2), we did not obtain a secure estimate of the $[OII]$ integrated flux lying outside of the slit, and thus we could not derive its $[OII]$ total luminosity. We roughly measured, though with large uncertainties, an observed $[OII]$ equivalent width $EW \approx 40 \text{ \AA}$, i.e., $\approx 20 \text{ \AA}$ in the rest frame.




FIG. 1.— *Top*: The dereddened VLT spectrum smoothed by 5 pixels (dotted line), overlaid with the template of a local Sc galaxy shifted to $z = 0.8424$ (solid line). *Bottom*: A zoom in the $\sim 6800 - 7300 \text{ \AA}$ range (dotted line), with the smoothed continuum (solid line) superimposed to emphasize on the possible detection of several stellar absorption lines as indicated in the panel.

In addition to the $[OII]$ emission doublet, we tentatively detect several stellar absorption features at a similar redshift such as $H12 (\lambda 3750 \text{ \AA})$, $H10 (\lambda 3798 \text{ \AA})$, $H9 (\lambda 3835 \text{ \AA})$ and $CaII K (\lambda 3934 \text{ \AA})$. The reliability of these features is yet questionable given the low signal to noise ratio in the continuum. Finally, in spite of the poor sky subtraction longward of 7300 \AA , there is a hint for a break in the continuum around 7400 \AA (see Figure 1). This suggests the presence of the rest-frame 4000 \AA -break as indicated by the comparison of our spectrum with the template of an Sc galaxy (Mannucci et al. 2001) shifted to $z = 0.84$, and provides further support to our redshift determination.

¹² see also http://www.ifa.au.dk/~hst/grb_hosts/data/grb990705/index.html

3.2. Structural parameters of the galaxy

High resolution HST images reveal that the host is a face-on Sc spiral galaxy. In Figure 2 we show a pseudo true-color image of that source, constructed by registering the deconvolved CL and LP data. Two primary spiral arms following a “ $m=2$ ” wave density mode clearly extend from the northern and southern sides of the central bulge, while other secondary arms are also observable. The disk spreads over a region of $\sim 3''$ in diameter, with a half-light radius $R_{0.5} \sim 7.5$ kpc ($0.9''$) in the range of those characterizing the disk-dominated galaxies at $0.75 \leq z \leq 1$ (Lilly et al. 1998).

After subtracting the foreground stars, we performed photometry within a $2''$ -radius aperture centered on the host nucleus using the STIS zero points of Gardner et al. (2000). Dereddened AB magnitudes $M_{AB}(\text{CL}) = 22.45 \pm 0.10$ and $M_{AB}(\text{LP}) = 22.0 \pm 0.1$ were respectively derived from the CL and LP images, the uncertainties being dominated by the subtraction residuals of the stars projected ahead of the galaxy.




FIG. 2.— A pseudo true-color image of the GRB 990705 host galaxy, resulting from the combination of the drizzled and deconvolved images taken through the CL and LP apertures. The 3σ -error location of GRB 990705 event as derived by Bloom et al. (2002) is indicated by the dashed circle. Note that the GRB occurred in the outskirts of a star forming region within one of the primary spiral arms. See the electronic edition of the *Journal for a color version of this figure*.

An attempt was made to fit the surface brightness profile of the galaxy using a “bulge + exponential disk” decomposition. Disentangling between the two contributions in distant sources requires not only a proper correction from the PSF effects (e.g., Moth02 but also a sufficient sensitivity up to ~ 10 to 15 kpc in the averaged profile of the disk to reliably constrain the scale length of the exponential component (see e.g., Fig. 1 of Rigopoulou02). Because of the diffuse and extended emission of the galaxy and the presence of the foreground stars, this was hardly achieved in our data. With large uncertainties, we suggest

however that the host galaxy is dominated by a large exponential disk (scale length ~ 5 – 6 kpc), with a rather small bulge contribution leading to a bulge-to-total light ratio $B/T \approx 0.10$ – 0.15 . We also estimated the central surface brightness of the disk component in the dereddened LP data $\mu_0(\text{LP})$, which was then converted into a rest-frame B_{AB} magnitude by applying a cosmological dimming term and a k -correction color factor as follows:

$$\mu_0(B_{AB}) = \mu_0(LP_{AB}) - 2.5 \log(1+z)^3 + (B_{obs} - LP)_{AB}$$

Since the LP aperture samples the almost rest-frame B emission at the redshift of the host, the color term should be rather small. Using the spectral energy distribution (SED) of a local Sc spiral galaxy shifted to $z = 0.84$ (see section 3.3), we derive a k -correction ~ 0.1 mag and finally obtain a value $\mu_0(B_{AB}) \approx 20.8$. This is actually less than the canonical Freeman (1970) value $\mu_0(B_{AB}) = 21.6$ observed in local disks, but is consistent with those found in higher- z spirals (Lilly et al. 1998). It is therefore in agreement with the observed global trend for the disk central surface brightness to significantly increase with redshift up to ~ 1 (Lilly et al. 1998).

3.3. Absolute B magnitude and star formation rate

At $z=0.84$, the rest-frame B emission of the galaxy is shifted to ~ 8100 Å. The dereddened continuum of the host of GRB 990705 and the CL – LP color derived from the STIS images are however consistent with the SED of a local Sc spiral galaxy (Mannucci et al. 2001) shifted to $z=0.84$ assuming no evolution. To estimate the luminosity at B_{rest} , we thus extrapolated the continuum of our spectrum using the template of Mannucci et al. (2001) and found a flux density $F_\nu(\lambda = 8100 \text{ Å}) \approx 12 \mu\text{Jy}$. Given the assumed cosmology and the luminosity distance of the galaxy, this implies a rest-frame absolute B-band magnitude $M_B \approx -21.75$ corresponding to a $2 L_*$ galaxy at $z \sim 1$ (Lilly et al. 1995).

Using a similar method, we also measured the continuum luminosity at $\lambda_{rest} = 2800$ Å to estimate the level of UV-unobscured star formation activity. From the Sc template SED, we estimate a flux $\sim 1.8 \mu\text{Jy}$ at the corresponding $\lambda_{obs} = 5152$ Å and deduce a UV luminosity $L_{UV} \sim 4 \text{ erg s}^{-1} \text{ Hz}^{-1}$ at the redshift of the host. Following the calibration of Madau et al. (1998) and assuming a Salpeter (1955) or Scalo (1986) Initial Mass Function, we finally derive star formation rates $\text{SFR} \approx 5 M_\odot \text{ yr}^{-1}$ or $\text{SFR} \approx 8 M_\odot \text{ yr}^{-1}$, which are fairly common values for star-forming galaxies at $z \sim 1$ (Lilly et al. 1995).

4. DISCUSSION AND CONCLUSION

The general properties of the GRB 990705 host have been summarized in Table 1. According to its morphology, star formation activity and absolute luminosity, we find that it is typical of the (disk) galaxies in the field at similar redshifts.

Taking account of the cumulative surface density distribution of sources with $R \leq 22.8$, Masetti et al. (2000) had estimated a probability of only 0.006 for the burst and the underlying galaxy being hazardously superimposed on the sky by projection effect, and had thus suggested a secure identification of this galaxy with the host of GRB 990705. Our spectroscopic observations reveal that

the redshift of the spiral is consistent with the one derived by Amati et al. (2000) for the GRB itself, providing further convincing evidence for a true association between the two. Among the current sample of GRB host galaxies, the host of GRB 990705 has been so far the only case clearly identified with a large disk-dominated spiral structure at high redshift, the others being classified either as compact, irregular or interacting systems (? , see e.g., Fig. 2 of) Bloom02a. With an absolute magnitude $M_B \approx -21.75$ ($H_0=65$, $\Omega_m=0.3$, $\Omega_\Lambda=0.7$), it lies furthermore within the brightest sources of the GRB host sample which is mostly characterized by sub-luminous systems.

With a larger sample of GRB hosts, the redshift-dependant proportion of large disks similar to the host of GRB 990705 relative to sub-luminous blue galaxies could provide indications on the fraction of star formation taking place in massive spirals and thus inform us of the cosmological evolution of the disk-dominated systems. This perspective appears promising since such massive and spiral objects are believed to be responsible for an important fraction of the Extragalactic Infrared Background (Rigopoulou et al. 2002). They could thus harbor star-forming regions enshrouded in dusty environments which are not sampled by the blue faint galaxy population.

We finally stress on the remarkable result obtained by Amati et al. (2000) who derived a reliable estimation of the burst redshift interpreting a transient edge observed in the GRB X-ray spectrum as an iron absorption at $z=0.86 \pm 0.17$. They showed that intrinsic GRB properties, such as the redshift and the physical conditions of the GRB-surrounding medium, can be derived from the burst detection itself, without the need of any afterglow to be detected and followed-up.

Eventhough GRB 990705 is the only one burst in which such a transient edge has been observed so far, which raises the question whether particular ionizing states of the circum-burst environment are required to detect these absorptions, this burst lies among the brightest GRBs ever detected with the Beppo-SAX satellite (Amati et al. 2000). This suggests that these transient edges could be

a more common feature of GRB spectra. Future satellites equipped with more sensitive X-ray detectors, such as the ECLAIRs experiment (Barret 2002), could be entirely dedicated for studying the GRB prompt emission, and may provide a systematic detection of these absorption lines. Larger samples of GRB redshifts could be derived, an achievement indeed required to estimate the star formation history in the Universe from the GRB occurrence rate.

Furthermore, compelling key results could be obtained towards the class of short GRBs or specific sub-classes of long GRBs such as the so-called dark bursts. The latter, exhibiting X-ray and radio afterglows without any detected optical counterparts, could pinpoint not only GRBs with optical afterglows either locally absorbed by dust or characterized by steep and rapid decays with time, but also very high redshift GRBs whose optical emission may be suppressed by the Gunn-Peterson HI trough along their line of sight. The use of transient features in X-ray spectra to derive GRB redshifts could thus provide a new approach to probe very distant GRBs in the early Universe. In the case of short GRBs, their distance scale and physical origin are simply still unknown since no detailed follow-up of their afterglows has been possible so far (? , but see) Castro-Tirado02. The clues of their formation mechanism directly observed in the GRB prompt emission would undoubtedly improve our current understanding of these particular events.

This work largely benefited from the input of public HST data taken as part of the "Survey of the Host Galaxies of Gamma-Ray Bursts" by Holland et al. (2000b). We wish to thank especially J. Greiner for his careful reading of the manuscript, as well as F. Masset, P. Goldoni and F. Daigne for fruitful discussions on this work. We also acknowledge the referee for his/her useful comments. We made extensive use of publicly available software with material credited to STScI and prepared for NASA under Contract NAS5-26555. This work was partially supported by CONICET/Argentina and Fundacion Antorchas.

REFERENCES

- Amati, L., Frontera, F., Vietri, M., et al. 2000, *Science*, 290, 953
 Barret, D. 2002, in *Proc. "GRB and Afterglow Astronomy 2001"* AIP Conf. Proc. in press (astro-ph/0205346)
 Blain, A. W. & Natarajan, P. 2000, *MNRAS*, 312, L35
 Bloom, J. S., Kulkarni, S. R., & Djorgovski, S. G. 2002, *AJ*, 123, 1111
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
 Castro-Tirado, A. J., Castro Ceron, J. M., Gorosabel, J., et al. 2002, *A&A*, accepted
 Djorgovski, S. G., Frail, D. A., Kulkarni, S. R., et al. 2001, *ApJ*, 562, 654
 Dutra, C. M., Bica, E., Clariá, J. J., Piatti, A. E., & Ahumada, A. V. 2001, *A&A*, 371, 895
 Freeman, K. C. 1970, *ApJ*, 160, 811
 Gardner, J. P., Baum, S. A., Brown, T. M., et al. 2000, *AJ*, 119, 486
 Holland, S., Andersen, M., Hjorth, J., et al. 2000a, *GCN* 753
 Holland, S., Fynbo, J., Thomsen, B., et al. 2000b, *GCN* 698
 Lamb, D. Q. & Reichart, D. E. 2000, *ApJ*, 536, 1
 Lazzati, D., Ghisellini, G., Amati, L., et al. 2001, *ApJ*, 556, 471
 Lilly, S., Schade, D., Ellis, R., et al. 1998, *ApJ*, 500, 75
 Lilly, S. J., Tresse, L., Hammer, F., Crampton, D., & Le Fevre, O. 1995, *ApJ*, 455, 108
 Madau, P., Pozzetti, L., & Dickinson, M. 1998, *ApJ*, 498, 106
 Mannucci, F., Basile, F., Poggianti, B. M., et al. 2001, *MNRAS*, 326, 745
 Masetti, N., Palazzi, E., Pian, E., et al. 2000, *A&A*, 354, 473
 Metzger, M. R., Djorgovski, S. G., Kulkarni, S. R., et al. 1997, *Nature*, 387, 878
 Mirabel, I. F., Sanders, D. B., & Le Floc'h, E. 2000, in "Cosmic Evolution and Galaxy Formation". ASP Conf. Series., Vol. 215, (astro-ph/0004022)
 Moth, P. & Elston, R. J. 2002, *AJ*, in press (Vol. 124, October 2002)
 Rigopoulou, D., Franceschini, A., Aussel, H., et al. 2002, *ApJ*, in press
 Salpeter, E. E. 1955, *ApJ*, 121, 161
 Saracco, P., Chincarini, G., Covino, S., et al. 2001, *GCN* 1010
 Scalo, J. M. 1986, *Fundamentals of Cosmic Physics*, 11, 1
 Starck, J. L., Murtagh, F., & Bijaoui, A. 1998, "Image Processing and Data Analysis", Cambridge University Press
 Totani, T. 1997, *ApJ*, 486, L71
 Vreeswijk, P. M., Fruchter, A., Kaper, L., et al. 2001, *ApJ*, 546, 672
 Wijers, R. A. M. J., Bloom, J. S., Bagla, J. S., & Natarajan, P. 1998, *MNRAS*, 294, L13
 Yoshida, A., Namiki, M., Otani, C., et al. 1999, *A&AS*, 138, 433

TABLE 1
PROPERTIES OF THE GRB 990705 HOST GALAXY

Parameter	Measure
α (J2000.0) (host nucleus)	$05^h 09^m 54^s.8$
δ (J2000.0) (host nucleus)	$-72^\circ 07' 54''$
Foreground extinction $E(B-V)$	0.12
Spectroscopic redshift	0.8424 ± 0.0002
CL aperture (dered.) magnitude ($\lambda_o = 5835 \text{ \AA}$)	22.45 ± 0.10
LP aperture (dered.) magnitude ($\lambda_o = 7208 \text{ \AA}$)	22.00 ± 0.10
Half-light radius $R_{0.5}$	$\sim 7.5 \text{ kpc}$
Central surface brightness $\mu_0(B_{AB})$	~ 20.8
Absolute M_B magnitude ($H_0=65, \Omega_m=0.3, \Omega_\lambda=0.7$)	≈ -21.75
<i>UV-unobscured</i> Star Formation Rate	$\sim 5\text{--}8 \text{ M}_\odot \text{ yr}^{-1}$